

## Assessment of the California mastitis test usage in smallholder dairy herds and risk of violative antimicrobial residues

Anakalo Shitandi\* and Gathoni Kihumbu<sup>1</sup>

Antibiotic Research Laboratory, Swedish University of Agricultural Sciences, Sweden

<sup>1</sup>Microbiology Laboratory, Faculty of Agriculture, Egerton University Njoro, Kenya

This study evaluated how predictive the California Mastitis Test (CMT) is for sub-clinical mastitis under tropical smallholder dairy production conditions in Kenya. It intended to establish whether the CMT usage could be contributing to misdiagnosis and consequent mistreatment with animal drugs resulting in residue problems. Milk samples (n = 239) were aseptically collected from lactating cows in the Rift Valley of Kenya and tested using the CMT, somatic cell counts (SCC) and bacterial culture. The samples were also screened for violative drug residues using the commercial delvo test and compared to the milks mastitic status for possible association. There was a numerical but non-significant ( $p > 0.05$ ) difference evident in the frequencies observed using the three different mastitis indicators. The prevalent bacterial species isolated from mammary glands with subclinical mastitis were *Staphylococcus aureus* (45.6%), coagulase-negative Staphylococci (13.0%), Streptococci (11.7%) and *Escherichia coli* 5.9%. There was an overall poor but significant ( $p < 0.05$ ) correlation between the CMT and the violative antimicrobial residues in samples from all quarters, infected and non-infected respectively. The results suggest that the CMT use amongst the smallholder dairy sector as a mastitic indicator may not be a risk factor in violative antimicrobial residues problems in milk.

**Key words:** California mastitis test, mastitic indicator, antimicrobial residues, milk

### Introduction

In lactating dairy animals the usage of antimicrobials is often aimed at treatment of mastitis an important diseases in

dairy herds [14]. The disease is usually classified into two forms, clinical in which the disease is diagnosed visually or by palpation, and subclinical which is mainly diagnosed via assessment of somatic cell and/or bacterial culture [1,2].

Veterinarians and milk producers use various mastitis indicators to help guide treatment decisions [5,4] although information on mastitis screening tests and their relevance for application in tropical conditions are scanty. The California Mastitis test, (CMT) is one such test utilized widely to determine the disease status of lactating animals, as it is simple, inexpensive and rapid screening test. The CMT reagent reacts with material from the nuclei of the somatic cells in the milk to form a gel and it estimates the number of somatic cells present in milk. The reaction is then visually scored, with interpretation dependant upon the amount of gel that forms [8].

In Kenya, many smallholder farmers being resource poor opt to use the California Mastitis Test (CMT) to detect sub-clinical mastitis infection. They do not take milk cultures to diagnostic laboratories, as there is widespread lack of diagnostic services due to the costs involved and limited laboratories. On the basis of a positive CMT score farmers then use antimicrobials for therapeutic treatment of the disease. The use of CMT is based on the increased number of somatic cells in milk and udder. There is however some evidence that the number of somatic cells in milk is relatively higher under tropical dairy production conditions, especially smallholder dairying [12]. This is attributable to poor hygienic conditions and high incidences of infections common in tropical dairy production systems. It was thus hypothesized that under smallholder dairy production conditions, positive sub-clinical mastitis based on CMT results may not be good indication of the presence of bacteria that cause udder infections. The study thus intended to establish whether the CMT usage could be contributing to misdiagnosis and consequent mistreatment with animal drugs resulting in the residue problems as evident from a previous study [19]. The research question of interest was how predictive the California mastitis test (CMT) is under smallholder dairy production conditions, for the presence of

\*Corresponding author

Phone: +46 18 67 20 60; Fax: + 46 18 67 29 95

E-mail: anakalo.shitandi@lmv.slu.se

pathogenic bacteria that cause sub-clinical mastitis.

## Materials and Methods

### Milk samples

A cross sectional study was done in the Rift Valley, a major milk production area of Kenya. Milk samples from 239 quarters (from 45 herds, which were crosses of Holstein Friesian and local breeds), were obtained by stratified random sampling. Milk samples were collected from the same herds at 1-month intervals until the calves were weaned. The samples were collected while the cows were restrained in a standing position. Udder and teats were cleaned with water and left to dry and then were swabbed with cotton soaked in 70% ethyl alcohol. The first three streams were discarded before 5 ml of milk were collected in sterile containers. Milk samples were initially screened with the CMT test reagent and then transported to the microbiological laboratory, Egerton University on ice and tested within 6 h of collection.

### CMT

The CMT reagent (DeLaval, Wroclaw, Poland) was used and the test was carried out using the method described by [17]. The reaction was then visually scored as 1, 2, 3, 4 or 5 according to the Scandinavian recommendations [7] depending upon the amount of gel that forms. The reaction was interpreted as follows: score 1 = no reaction; score 2 = slight slime which disappears with continued swirling; score 3 = distinct slime but without gel formation; score 4 = immediate formation of gel which moves as a mass during swirling; score 5 = gel develops a convex surface and adheres to the bottom of the paddle.

### Somatic cell count

Somatic cell counts (SCC) were determined by spreading 10  $\mu$ l of thoroughly mixed milk from each sample over 1 cm<sup>2</sup> area on a glass slide. The slides were left to air-dry and were stained by Newman-Lampert stain as described by [6]. SCCs of  $\geq 10^5$  cell/ml milk were considered to be positive.

### Bacteriology

Isolation and identification were carried out according to the Scandinavian recommendations on examination of quarter milk samples [9]. In brief, immediately after delivery, the milk samples were inoculated on blood agar plates (Becton Dickinson and Company, USA), which were divided into four sections. A disposable culturing loop (10  $\mu$ l) was dipped into the milk sample and a total of 6-8 lines made in one agar section by turning the loop once between the streaking lines. Each quarter milk sample (10  $\mu$ l) was thus streaked in one section of the plate as a line culture. Samples were cultivated under aerobic conditions

for 24-72 h at 37°C and examined for bacterial growth. Pure cultures were further examined for morphological, staining and cultural characteristics, and for biochemical reactions. In cases of mixed growth, a new quarter sample was taken and reexamined. Where no growth was detected, plates were reincubated at 37°C for an additional 24 h. Bacteria were identified using a standardized procedure [10]. API 20 Strep and API Staph (BioMérieux, Basingstoke, Hants) were used for speciation of Streptococci and coagulase-negative *Staphylococcus*. When fungal infection was suspected, portions of the milk were cultured on Sabouraud dextrose agar (Difco, Detroit, MI, USA).

Quarters were classified as not infected if no organisms were isolated and infected with major pathogens if *Streptococcus agalactiae*, other *Streptococcus* species, *Staphylococcus aureus*, and *Escherichia coli*, coagulase-negative *staphylococci* were isolated. Quarters were also classified as “normal” if no organisms were isolated, the udder had no injuries or indurations, the appearance of the milk was normal, and no previous history of mastitis was recorded and “abnormal” otherwise. Samples with unspecified mixed cultures were considered contaminated and thus excluded from subsequent analysis.

### Violative antimicrobial residues

All samples were examined the same day for the presence of antimicrobial residues. The milk samples were all preheated at were pre-heated at 80°C for 10 min to inactivate natural inhibitory substances and kill contaminating bacteria. The commercial Delvotest SP (GistBrocades, Delft, the Netherlands) test was then used to screen the samples which was performed as per the manufacturer and included appropriate positive and negative control samples in all assays. The test is a microbial inhibitor test, which utilises *Bacillus stearothermophilus* as the test organism with a dye in the media and detects antimicrobial substances non-specifically. Under normal conditions as the culture grows the dye color is changed from purple to yellow. If an antibiotic is present the organism is inhibited and the dye remains purple.

### Statistical analysis

For comparison of the methods (SCC, CMT and bacteriological), statistically significant association was determined by the Chi-squared test, based on cut off for results as positive or negative. All statistical analysis was performed using the Minitab, 2000 program [9]. Statistical significance was set at the 5% level. Data was assessed by least square analysis of variance, using the general linear model procedure. The model included the effects of the animal, the quarter, the stage of lactation (1-12 months), the bacteriological findings and the antimicrobial status. The results are presented as means  $\pm$  SEM. The correlation coefficient between the mastitic indicator-CMT and

**Table 1.** Observed frequencies (%) of milk samples (n = 239) positive and negative for mastitis using the CMT, SCC, and bacteriological mastitic indicators from apparently unaffected cows

	CMT	SCC <sup>a</sup>	Bacteriological
Negative	39 (16.3%)	48 (20.1%)	57 (23.8%)
Positive	200 (83.7%)	191 (79.9%)	182 (76.2%)

<sup>a</sup>SCCs of >10<sup>5</sup> cell/ml milk was considered positive  
P value > 0.05

**Table 2.** Bacteriological findings (%) in quarter samples (n = 239) from smallholder cows

Infection Status	Proportion N (%)				
	Non infected	<i>Staphylococcus aureus</i>	Coagulase negative staphylococci	<i>Streptococci spp</i>	<i>Escherichia coli</i>
Proportion (%)	57 (23.8)	109 (45.6)	31 (13.0)	28 (11.7)	14 (5.9)

**Table 3.** †Mean ± SEM of SCC and CMT for infected and non infected quarters of cows

Component	n	Infected (×10 <sup>5</sup> )	Non infected (×10 <sup>3</sup> )	P-value
SCC	186	18.24 ± 0.24 (n = 148)	20.05 ± 0.39 (n = 38)	p < 0.01
CMT score	239	1.98 ± 0.14 (n = 18)	23.49 ± 0.29 (n = 57)	p < 0.001

†The data are presented as mean cells per milliliter ± standard error of the mean for cows in each group.  
SCC = somatic cell count; CMT = California mastitis test; n = number of observations.

**Table 4.** †Mean ± SEM of SCC and CMT in non -infected and in *S. aureus* and CNS -infected quarters of cows

Component	N	<i>S. aureus</i> (×10 <sup>5</sup> )	Non infected (×10 <sup>3</sup> )	CNS
SCC	186	18.24 ± 0.24 (n = 109)	19.53 ± 0.59 (n = 46)	19.47 ± 0.89 (n = 31)
CMT score	209	1.98 ± 0.14 (n = 105)	3.11 ± 0.36 (n = 81)	3.38 ± 0.30 (n = 13)

†The data are presented as mean cells per milliliter ± standard error of the mean for cows in each group.  
SCC = somatic cell count; CMT = California mastitis test; CNS = coagulase negative staphylococci  
n = number of observations.

violative antimicrobial residues (as measured by the Delvo test SP) was calculated using Spearman rank correlation of the residual obtained after correcting for the effects in the statistical model.

## Results

Observed frequencies of milk samples from apparently unaffected cows, which were positive and negative for mastitis using the three mastitic indicators (CMT, SCC and bacteriological), are displayed in Table 1. While there was a numerical difference evident in the frequencies observed using the mastitis indicators the difference was not significant (p value = 0.121 > 0.05).

From Table 2, intramammary infections were present in 76.2% of the 239-quarter milk samples examined. *S. aureus* and Coagulase negative staphylococci (CNS) represented 59.9% and 17.0% of the isolates respectively with *Streptococci* representing 15.4%.

The mean values for the SCC and CMT from the infected and non-infected quarters are shown in Table 3. Infected

udder quarters had significantly (p < 0.01) higher mean values for both SCC and CMT. Eleven of the 239 individual quarter samples had somatic counts >5,000,000/ml which was considered very elevated for apparently normal milk. There were also eight samples of the 239 individual quarter samples that showed high values for CMT and SCC from which bacteria were not isolated.

Table 4, shows separately the mean values for non-infected quarters and quarters infected by CNS and *S. aureus*. From the statistical model, the correlation coefficient between the mastitic indicator-CMT and violative antimicrobial residues (as measured by the Delvo test SP) was calculated using Spearman rank correlation. The correlations from all quarters was 0.59 (n = 186); infected quarters had a correlation of 0.38 (n = 148) and non-infected was 0.48 (n = 38). All positive and negative control samples yielded the expected reactions and >12% of the milk samples collected from the herds contained inhibitory substances which were heat stable. There was an overall poor but significant (p < 0.001) correlation between the CMT and the violative antimicrobial residues.

## Discussion

The relationship between the mastitis indicators, CMT, SCC and bacterial culture was investigated based on the proportions of positives at set cut off limits. The null hypothesis was that the true mastitis rates are the same for the three indicators with the alternative hypothesis being that they are not equal. The results suggest that under smallholder dairy production conditions, positive sub-clinical mastitis based on CMT results may be good indication of the presence of bacteria that cause udder infections. The CMT may thus not be contributing to misdiagnosis and consequent mistreatment with animal drugs resulting in the residue problems.

While the spearman's rank correlation coefficient provided a measure of association between the two variables it does however not imply causality, although in some studies [15,16] an association has been observed between measures of milk quality and risk of violative antimicrobial residues.

There are however many factors that contribute to the overuse of antimicrobials in farming which arises from a complex interaction between the cow, microbial agents, environmental influences and management factors [18]. The Delvotest microbial inhibitor test can be inhibited by lysozyme or lactoferrin separately and synergistically [3]. These inhibitors and other non-protein inhibitors occur frequently in high SCC (>4,000,000/ml) milk and could be a problem in tropical conditions where SCC has been observed to be abnormally high [12], a finding also observed in this study. The inhibitors are however heat labile and hence the need to heat-treat the samples at >80°C as was done in this study.

A critical aspect in the evaluation of screening tests is the criteria of deciding upon the true status of the udder. In previous studies, bacteriological findings, SCC, or a combination has been used [5,7]. *S. aureus* is regarded in many regions as a major causative organism for mastitis [11,13] and this was found to be the case in the study site justifying its possible use as a true status of udder infection within the region.

The field study provided no clear evidence that CMT could be less predictive compared to the other two mastitis indicators in the small herd sector. The test is thus unlikely to be contributing to misdiagnosis and consequent mistreatment with animal drugs resulting in the violative residue problems inherent in milk. There were however samples that showed high values for CMT and SCC from which bacteria were not isolated and this together with the apparently high rate of intramammary infection in the smallholder dairy sector warrants further investigation. In the model that was used to determine possible association between CMT usage violative residue problems although the observed correlation was weak, the observed frequencies

of antimicrobial residues emphasize the importance of screening farm milk for antibiotic residues.

In conclusion the results suggest that the CMT may be a useful indicator of udder infection on farm that should however be used alongside bacteriological culture. A program is also needed in the region, designed to improve milk quality and udder health so as to prevent new intramammary infections, eliminate existing infections and monitor udder health status.

## Acknowledgments

We are grateful for financial support from the Swedish Institute (SI) - Sweden. We thank Dr. Sternesjö, Åse (SLU, Uppsala) and Dr. Carlos Concha (Swedish Veterinary Association, Uppsala) for their material assistance and acknowledge the input of Emil Olden, Claes Ericksson (SLU) in sample collection and laboratory analysis.

## References

- Blood D, Henderson J, Rodostits M.** Veterinary Medicine, pp. 501-550. 8th ed. ELBS. London.1990.
- Blowey R, Edmondson P.** Mastitis Control in Dairy Herds. pp. 77-89. Farming Press, Ipswich, UK. 1995.
- Carlsson Å, Björck L.** The effect of some indigenous antibacterial factors in milk on the growth of *Bacillus stearothermophilus* var. *calidolactis*. *Milchwissenschaft*. 1987, **42**, 282-285.
- Deluyker H, Gay J, Weaver A.** Interrelationships of somatic cell count, mastitis and milk yield in a low somatic cell count herd. *J Dairy Sci* 1993, **76**, 3445-3452
- Emmanuelson U, Olsson T, Holmberg O, Hageltorn M, Mattila T, Nelsson L, Astrom G.** Comparison of some screening test for detecting mastitis. *J Dairy Sci* 1987, **70**, 880-887.
- Emmert H.** Veterinary Clinical Pathology. 4th ed, pp. 34-39, Saunders, Philadelphia, 1986.
- Grootenhuis G, Oldenbroek J, Van den Berg.** Differences in mastitis susceptibility between Holstein Friesian, Dutch Friesian and Dutch Red and White cows. *Vet Quart* 1979, **1**, 37-39.
- Klastrup O, Schmidt P.** Nordic recommendations concerning mastitis control of quarter samples. *Nordic Veterinarian-Medicine*, 1974, **26**, 197-204.
- Minitab statistical software.** For windows 2000. Mintab Inc. USA, 2000.
- National Mastitis Council.** Laboratory Handbook on Bovine Mastitis. pp. 22-47. National Mastitis Council, Inc. 2820 Madison. USA, 1999.
- Nickerson S, Owens W, Fox K, Scheifinger C, Shryock R, Spike E.** Comparison of tilmicosin and cephalixin as therapeutics for *Staphylococcus aureus* mastitis at dry-off. *J Dairy Sci* 1999, **82**, 696-699.
- Omoro A.** Epidemiology and economics of mastitis in the smallholder dairy sector of Kiambu District, Kenya. pp. 14-

18. PhD. Thesis, University of Nairobi. 1997.
- 13. Osteras O, Edge V, Martin S.** Determinants of success or failure in the elimination of major mastitis pathogens in selective dry cow therapy. *J Dairy Sci* 1999, **82**, 122-125.
- 14. Radostits O, Blood D, Gay C** (ed.). *Veterinary Medicine. A textbook of the Diseases of Cattle, Sheep, Pigs, Goats and Horses.* pp. 48-52. Saunders, Philadelphia, 2000.
- 15. Ruegg P, Tabone J.** The relationship between antibiotic residues violations and somatic cell counts in Wisconsin dairy herds. *J Dairy Sci* 2000, **83**, 2805-2809.
- 16. Saville J, Wittum T, Smith K.** Association between measures of milk quality and risk of violative antimicrobial residues in grade-A raw milk. *J Am Vet Med Assoc* 2000, **217**, 541-545.
- 17. Schalm O, Noorlander O.** Experiments and observations leading to development of California Mastitis Test. *J Am Vet Med Assoc* 1957, **130**, 199-204.
- 18. Seers P, Gonzalez R, Wilson J, Han R.** Procedures for mastitis diagnosis and control. *Vet Clin North Am Food Anim Pract* 1993, **9**, 445-468.
- 19. Shitandi A, Sternesjö A.** Detection of antimicrobial drug residues in Kenyan milk. *J Food Prot* 2001, **21**, 205-214.

The California mastitis test (CMT) is a simple cow-side indicator of the somatic cell count of milk. It operates by disrupting the cell membrane of any cells present in the milk sample, allowing the DNA in those cells to react with the test reagent, forming a gel. It provides a useful technique for detecting subclinical cases of mastitis. Whiteside described a reaction between sodium hydroxide and milk that resulted in the thickening of mastitic milk. The utility of this reaction as a field test was The antimicrobial use assessment tool was developed and set up in Open Data Kit (ODK) on mobile tablet devices. The tool included open-ended and closed questions about household demographics, farm characteristics, management of manure, feed types, animal health constraints, disease prevention, animal health services, antimicrobial use, animal product consumption, and costs related to animal health. Prior to the study, veterinarians in the localities were trained as enumerators and the questionnaire was piloted with 40 livestock owners as a first step of validating the tool. IRT analysis, which provides information on the discrimination and difficulty of each item across different levels of the underlying trait, was used. Incidence of dairy cow mastitis and associated risk factors in sodo town and its surroundings , wolaita zone , ethiopia. A. Asmare, F. KASSA Wolaita2017. Evaluation of vaccine against Escherichia coli mastitis in dairy cattle. E. El-Ebiary, H. M. Ibrahim, Makharita2014. Assessment of the California mastitis test usage in smallholder dairy herds and risk of violative antimicrobial residues. Mastitis is an increasing problem in dairy farms and in the dairy industry. The paper presents changes in susceptibility of strains of bacteria isolated from quarter milk cows from mastitis in 2011, 2012, 2013, 2014, 2015 in the Factory Laboratory of Dairy Cooperative in south-eastern Poland. In total, 3409 strains of staphylococci, 2774 strains of streptococci and 423 strains of E. coli were isolated in these years. According to estimates in Poland, the clinical form of mastitis affects 6.5% of the udder quarters, while the sub-clinical form was observed in as many as 43.4% of the cases studied [18]. The problem of mastitis affects many European countries, where milk production is one of the key sectors of agriculture.